

Mechanisms of Plant Stress Tolerance: Drought, Salinity, and Temperature Extremes

Ritik Chawla^{1*}, Kinjal Mondal², Pankaj³, Meenakshi Sahu⁴

¹Department of Fruit Science, College of Horticulture, Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, 173 230-India.

²Department of Molecular Biology and Biotechnology, Maharana Pratap University of Agriculture and Technology Udaipur-313001, Rajasthan, India.

³Department of Botany and Plant Physiology, CCSHAU, Hisar, Haryana, India.

⁴Institute of agricultural Sciences, Bundelkhand University, Jhansi, Uttar Pradesh 284128, India.

Citation: Ritik Chawla, Kinjal Mondal, Pankaj, Meenakshi Sahu (2022). Mechanisms of Plant Stress Tolerance: Drought, Salinity, and Temperature Extremes. *Plant Science Archives*. 04-08. DOI: <https://doi.org/10.51470/PSA.2022.7.2.04>

Corresponding Author: Ritik Chawla | E-Mail: (ritikchawla0416@gmail.com)

Received 23 February 2022 | Revised 19 March 2022 | Accepted 11 June 2022 | Available Online 12 June 2022

ABSTRACT

Plants are continuously exposed to various environmental stresses that significantly affect their growth, development, and productivity. Among these, drought, salinity, and temperature extremes are the most detrimental. Understanding the mechanisms underlying plant stress tolerance is crucial for developing strategies to improve crop resilience and ensure food security. This review comprehensively explores the physiological, biochemical, and molecular mechanisms of plant tolerance to drought, salinity, and temperature extremes. We discuss the roles of stress perception and signaling, osmotic adjustment, antioxidant defense, hormonal regulation, and genetic and epigenetic modifications. Additionally, we highlight recent advancements in breeding and biotechnological approaches aimed at enhancing stress tolerance in crops.

Keywords: Signaling, osmotic adjustment, antioxidant defense, hormonal regulation

1. Introduction

Plants, as immobile organisms, face a variety of environmental challenges that can severely affect their growth, development, and productivity. Among these challenges, abiotic stresses such as drought, salinity, and temperature extremes (both high and low) are particularly detrimental. These stresses are becoming more frequent and intense due to climate change, posing significant threats to global agriculture and food security. Understanding the complex mechanisms underlying plant stress tolerance is critical for developing strategies to enhance crop resilience and ensure sustainable agricultural production in the face of these challenges [1-2].

Drought stress, characterized by a lack of adequate water availability, affects plant water relations, leading to reduced cell turgor, impaired photosynthesis, and stunted growth. Salinity stress, resulting from high concentrations of soluble salts in the soil, disrupts ion homeostasis and osmotic balance, causing ionic toxicity and physiological drought. Temperature extremes, including both heat and cold stress, can cause protein denaturation, membrane instability, and metabolic imbalances, further challenging plant survival and productivity [3-4]. Plants have evolved a range of adaptive mechanisms to cope with these stresses. These mechanisms operate at various levels, from cellular and molecular to physiological and biochemical. At the cellular level, stress perception and signaling pathways are activated in response to environmental cues, leading to the expression of stress-responsive genes and proteins. Physiologically, plants adjust their water status through osmotic adjustment and maintain ion homeostasis under saline conditions. Biochemically, antioxidant defense systems mitigate oxidative damage caused by reactive oxygen species (ROS) generated during stress. Hormonal regulation also plays a vital role in modulating stress responses, with hormones such as

abscisic acid (ABA), ethylene, and salicylic acid (SA) orchestrating complex signaling networks [5-6]. In addition to these intrinsic mechanisms, advances in breeding and biotechnological approaches have significantly contributed to enhancing plant stress tolerance. Traditional breeding methods, marker-assisted selection (MAS), and genomic selection (GS) have been employed to develop stress-tolerant crop varieties. Modern biotechnological tools, including genetic engineering and gene editing technologies like CRISPR/Cas9, allow for precise manipulation of stress-responsive genes, offering new avenues for improving crop resilience [7]. This review aims to provide a comprehensive overview of the mechanisms of plant stress tolerance to drought, salinity, and temperature extremes. We will explore the physiological, biochemical, and molecular responses of plants to these stresses, highlighting key components and processes involved in stress perception, signaling, osmotic adjustment, antioxidant defense, and hormonal regulation. Furthermore, discuss the role of genetic and epigenetic modifications in enhancing stress tolerance and examine recent advancements in breeding and biotechnological approaches [8]. By integrating insights from fundamental research with practical applications, this review seeks to contribute to the development of strategies for improving crop resilience and ensuring food security in the face of changing environmental conditions.

2. Drought Stress Tolerance

2.1. Stress Perception and Signaling

Plants perceive drought stress through changes in cell turgor, membrane fluidity, and osmotic pressure. These changes are sensed by specific receptors and ion channels, leading to the activation of signaling cascades involving calcium ions (Ca²⁺), reactive oxygen species (ROS), and various protein kinases.

Important signaling pathways include the mitogen-activated protein kinase (MAPK) pathway and the abscisic acid (ABA)-dependent pathway.

2.2. Osmotic Adjustment

Osmotic adjustment is a crucial mechanism that allows plants to maintain cell turgor under drought conditions. This involves the accumulation of compatible solutes, such as proline, glycine betaine, and sugars, which help to stabilize cellular structures and protect against dehydration [9].

2.3. Antioxidant Defense

Drought stress leads to the overproduction of ROS, which can cause oxidative damage to cellular components. Plants counteract this by enhancing their antioxidant defense systems, including enzymatic antioxidants like superoxide dismutase (SOD), catalase (CAT), and peroxidases (POD), as well as non-enzymatic antioxidants like ascorbate and glutathione [10].

2.4. Hormonal Regulation

Hormones play a vital role in regulating plant responses to drought stress. ABA is the primary hormone involved, promoting stomatal closure to reduce water loss and inducing the expression of drought-responsive genes. Other hormones, such as ethylene, salicylic acid (SA), and jasmonic acid (JA), also modulate drought tolerance by influencing various physiological processes [11].

3. Salinity Stress Tolerance

3.1. Ion Homeostasis

Salinity stress primarily affects plants through ionic toxicity and osmotic stress. Plants maintain ion homeostasis by regulating the uptake, transport, and compartmentalization of ions. The Na⁺/H⁺ antiporter SOS1 (Salt Overly Sensitive 1) and vacuolar Na⁺/H⁺ exchangers (NHX) are key components in maintaining low cytosolic Na⁺ concentrations [12].

3.2. Osmotic Adjustment

Similar to drought stress, osmotic adjustment plays a significant role in salinity tolerance. The accumulation of compatible solutes helps to maintain osmotic balance and protect cellular structures from the harmful effects of high salt concentrations [13].

3.3. Antioxidant Defense

Salinity stress also leads to oxidative stress, necessitating the activation of antioxidant defense mechanisms. The enhanced production of enzymatic and non-enzymatic antioxidants helps to mitigate the damaging effects of ROS [14].

3.4. Hormonal Regulation

Salinity stress induces changes in hormone levels and signaling pathways. ABA, ethylene, and cytokinins are particularly important in modulating plant responses to salinity. ABA enhances osmotic adjustment and antioxidant defense, while ethylene and cytokinins influence ion transport and stress-responsive gene expression.[15].

4. Temperature Stress Tolerance

4.1. Heat Stress Tolerance

4.1.1. Heat Shock Proteins

Heat stress leads to protein denaturation and aggregation, which can impair cellular functions. Heat shock proteins (HSPs) act as molecular chaperones, helping to refold damaged proteins and prevent aggregation. The expression of HSPs is regulated by heat shock factors (HSFs) [16].

4.1.2. Membrane Stability

Maintaining membrane stability is crucial for heat tolerance. Plants achieve this by altering the composition of membrane lipids, increasing the proportion of saturated fatty acids, and stabilizing membrane proteins.

4.1.3. Antioxidant Defense

Heat stress also induces oxidative stress, necessitating the activation of antioxidant defense systems. Enhanced production of antioxidants helps to protect cellular components from heat-induced oxidative damage.

4.1.4. Hormonal Regulation

Hormones such as ABA, ethylene, and brassinosteroids play significant roles in modulating plant responses to heat stress. ABA induces the expression of heat-responsive genes, while ethylene and brassinosteroids influence membrane stability and antioxidant defense.

4.2. Cold Stress Tolerance

4.2.1. Cold Acclimation

Cold acclimation is a process whereby plants increase their tolerance to low temperatures through gradual exposure. This involves changes in gene expression, protein function, and metabolic pathways.

4.2.2. Membrane Fluidity

Maintaining membrane fluidity is essential for cold tolerance. Plants adjust the composition of membrane lipids, increasing the proportion of unsaturated fatty acids to prevent membrane rigidity.

4.2.3. Antioxidant Defense

Cold stress induces the production of ROS, requiring the activation of antioxidant defense mechanisms. Enhanced production of antioxidants helps to mitigate oxidative damage under cold conditions [17].

4.2.4. Hormonal Regulation

Hormones such as ABA, gibberellins, and cytokinins are involved in regulating plant responses to cold stress. ABA promotes the expression of cold-responsive genes, while gibberellins and cytokinins influence growth and development under low temperatures.

Table 1: This table summarizes the mechanisms and components involved in plant stress tolerance, along with examples of breeding and biotechnological approaches aimed at enhancing crop resilience to drought, salinity, and temperature extremes.

| Stress Type | Mechanisms | Key Components/Processes | Examples of Breeding/Biotechnological Approaches |
|-----------------|---------------------------------|---|---|
| Drought | Stress Perception and Signaling | Receptors, ion channels, Ca ²⁺ , ROS, MAPK pathway, ABA-dependent pathway | Marker-assisted selection (MAS), CRISPR/Cas9, transgenic plants expressing DREB genes |
| | Osmotic Adjustment | Accumulation of compatible solutes (proline, glycine betaine, sugars) | Development of drought-tolerant maize varieties |
| | Antioxidant Defense | Enzymatic (SOD, CAT, POD) and non-enzymatic antioxidants (ascorbate, glutathione) | Genetic engineering for enhanced antioxidant production |
| | Hormonal Regulation | ABA, ethylene, salicylic acid (SA), jasmonic acid (JA) | Breeding for enhanced ABA response |
| Salinity | Ion Homeostasis | Na ⁺ /H ⁺ antiporters (SOS1), vacuolar Na ⁺ /H ⁺ exchangers (NHX) | Development of salinity-tolerant rice varieties using gene editing |
| | Osmotic Adjustment | Accumulation of compatible solutes | Breeding and biotechnological approaches to enhance osmotic adjustment |
| | Antioxidant Defense | Enzymatic and non-enzymatic antioxidants | Transgenic approaches for improved antioxidant defense |
| | Hormonal Regulation | ABA, ethylene, cytokinins | Genetic modification to enhance hormone regulation under salinity stress |
| Heat | Heat Shock Proteins | HSPs, heat shock factors (HSFs) | Transgenic plants overexpressing HSPs |
| | Membrane Stability | Alteration of membrane lipid composition | Breeding for membrane stability |
| | Antioxidant Defense | Enzymatic and non-enzymatic antioxidants | Gene editing for enhanced antioxidant defense |
| | Hormonal Regulation | ABA, ethylene, brassinosteroids | Breeding for enhanced hormonal responses |
| Cold | Cold Acclimation | Changes in gene expression, protein function, metabolic pathways | Development of cold-tolerant varieties through MAS and GS |
| | Membrane Fluidity | Increase in unsaturated fatty acids | Breeding for altered membrane lipid composition |
| | Antioxidant Defense | Enzymatic and non-enzymatic antioxidants | Genetic engineering for improved cold stress tolerance |
| | Hormonal Regulation | ABA, gibberellins, cytokinins | Breeding for hormonal regulation under cold stress |
| General | Genetic Modifications | Conventional breeding, transgenic plants, CRISPR/Cas9 | Integrated approaches combining conventional and modern techniques |
| | Epigenetic Modifications | DNA methylation, histone modifications, small RNAs | Epigenetic breeding for heritable stress tolerance |
| | Integrated Approaches | Phenotyping and genotyping integration, multiple stress tolerance traits | Comprehensive breeding programs utilizing genetic diversity and advanced tools |

5. Genetic and Epigenetic Modifications

5.1. Genetic Modifications

Genetic modifications, through conventional breeding or modern biotechnological approaches, have been used to enhance stress tolerance in crops. Transgenic plants expressing stress-responsive genes, such as DREB (dehydration-responsive element-binding) and HSPs, have shown improved tolerance to drought, salinity, and temperature extremes.

5.2. Epigenetic Modifications

Epigenetic modifications, including DNA methylation, histone modifications, and small RNAs, play crucial roles in regulating gene expression in response to stress. These modifications can be stable and heritable, providing a mechanism for plants to "remember" previous stress exposures and enhance tolerance in subsequent generations.

6. Breeding and Biotechnological Approaches

6.1. Conventional Breeding

Conventional breeding involves selecting and cross-breeding plants with desirable traits. Marker-assisted selection (MAS) and genomic selection (GS) have accelerated the breeding process by enabling precise selection at the DNA level.

6.2. Genetic Engineering

Genetic engineering, including transgenic and gene-editing technologies like CRISPR/Cas9, allows for the precise modification of stress-responsive genes. This has led to the development of crops with enhanced tolerance to drought, salinity, and temperature extremes.

6.3. Integrated Approaches

Combining conventional breeding with biotechnological

approaches offers a comprehensive strategy for developing stress-tolerant crops. Integrating phenotyping and genotyping data, utilizing genetic diversity, and targeting multiple stress tolerance traits are crucial for the success of these approaches.

7. Case Studies and Success Stories

7.1. Drought-Tolerant Maize

Drought-tolerant maize varieties, developed through conventional and molecular breeding, have been widely adopted in sub-Saharan Africa. These varieties show significant yield improvements under water-limited conditions [18].

7.2. Salinity-Tolerant Rice

Gene-edited rice varieties with enhanced salinity tolerance have shown promise in regions affected by soil salinization. These varieties exhibit improved growth and yield under high salt conditions.

7.3. Heat-Tolerant Wheat

Heat-tolerant wheat varieties, developed through marker-assisted breeding, have been successfully cultivated in regions prone to heatwaves. These varieties maintain yield and quality under high-temperature conditions [19-20].

8. Future Directions and Policy Implications

8.1. Research and Development

Continued research and development are essential for understanding the complex mechanisms of stress tolerance and developing new strategies to enhance crop resilience. Collaborative efforts between research institutions, governments, and the private sector are crucial for success.

8.2. Policy Support

Supportive policies that promote investment in advanced breeding technologies, capacity building for breeders, and the conservation of genetic diversity are essential. Regulatory frameworks that facilitate the adoption of genetically modified and gene-edited crops will also play a vital role.

8.3. Farmer Adoption

Ensuring that stress-tolerant crop varieties are accessible and affordable for farmers is critical. Extension services and training programs can help farmers adopt and implement these technologies, improving their resilience to environmental stresses [21-22].

9. Conclusion

Understanding the mechanisms of plant stress tolerance is fundamental for developing strategies to enhance crop resilience to drought, salinity, and temperature extremes. Advances in physiological, biochemical, and molecular research, combined with innovative breeding and biotechnological approaches, offer promising solutions to this challenge. By integrating genetic diversity, phenotyping, and genotyping, and focusing on multiple stress tolerances, we can safeguard global food systems and ensure agricultural sustainability in the face of climate change. Continued research, investment, and supportive policies are essential to realize these goals and secure the future of agriculture.

References

- Zhu, J.-K. (2001). Plant salt tolerance. *Trends in Plant Science*, 6(2), 66-71.
- Tardieu, F., & Tuberosa, R. (2010). Dissection and modelling of abiotic stress tolerance in plants. *Current Opinion in Plant Biology*, 13(2), 206-212.
- Xiong, L., Schumaker, K. S., & Zhu, J.-K. (2002). Cell signaling during cold, drought, and salt stress. *The Plant Cell*, 14(suppl_1), S165-S183.
- Mittler, R. (2002). Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science*, 7(9), 405-410.
- Verslues, P. E., Agarwal, M., Katiyar-Agarwal, S., Zhu, J., & Zhu, J.-K. (2006). Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *The Plant Journal*, 45(4), 523-539.
- Blum, A. (2017). Osmotic adjustment is a prime drought stress adaptive engine in support of plant production. *Plant, Cell & Environment*, 40(1), 4-10.
- Valliyodan, B., Nguyen, H. T. (2006). Understanding regulatory networks and engineering for enhanced drought tolerance in plants. *Current Opinion in Plant Biology*, 9(2), 189-195.
- Mahajan, S., & Tuteja, N. (2005). Cold, salinity and drought stresses: An overview. *Archives of Biochemistry and Biophysics*, 444(2), 139-158.
- Bray, E. A. (1997). Plant responses to water deficit. *Trends in Plant Science*, 2(2), 48-54.
- Wang, W., Vinocur, B., & Altman, A. (2003). Plant responses to drought, salinity and extreme temperatures: Towards genetic engineering for stress tolerance. *Planta*, 218(1), 1-14.
- Munns, R., & Tester, M. (2008). Mechanisms of salinity tolerance. *Annual Review of Plant Biology*, 59, 651-681.
- Chaves, M. M., Maroco, J. P., & Pereira, J. S. (2003). Understanding plant responses to drought - from genes to the whole plant. *Functional Plant Biology*, 30(3), 239-264.
- Flowers, T. J. (2004). Improving crop salt tolerance. *Journal of Experimental Botany*, 55(396), 307-319.
- Parida, A. K., & Das, A. B. (2005). Salt tolerance and salinity effects on plants: A review. *Ecotoxicology and Environmental Safety*, 60(3), 324-349.
- Yamaguchi-Shinozaki, K., & Shinozaki, K. (2006). Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annual Review of Plant Biology*, 57, 781-803.

16. Hirayama, T., & Shinozaki, K. (2010). Research on plant abiotic stress responses in the post-genome era: Past, present and future. *The Plant Journal*, 61(6), 1041-1052.
17. Reddy, A. R., Chaitanya, K. V., & Vivekanandan, M. (2004). Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of Plant Physiology*, 161(11), 1189-1202.
18. Ahmad, P., & Prasad, M. N. V. (Eds.). (2011). *Abiotic Stress Responses in Plants: Metabolism, Productivity and Sustainability*. Springer.
19. Bohnert, H. J., Gong, Q., Li, P., & Ma, S. (2006). Unraveling abiotic stress tolerance mechanisms - getting genomics going. *Current Opinion in Plant Biology*, 9(2), 180-188.
20. Haak, D. C., Fukao, T., Grene, R., Hua, Z., Ivanov, R., Perrella, G., & Li, S. (2017). Multilevel regulation of abiotic stress responses in plants. *Frontiers in Plant Science*, 8, 1564.
21. Knight, H., & Knight, M. R. (2001). Abiotic stress signalling pathways: Specificity and cross-talk. *Trends in Plant Science*, 6(6), 262-267.
22. Zhang, H., Zhang, J., Lang, Z., Botella, J. R., & Zhu, J.-K. (2017). Genome editing—principles and applications for functional genomics research and crop improvement. *Critical Reviews in Plant Sciences*, 36(4), 291-309.