

An Overview of Mycorrhiza in Pines: Research, Species, and Applications

Bharathiraja¹ and Lalit Upadhyay^{2*}

¹Department of Agriculture Microbiology, Agriculture College and Research Institute, TNAU-Chettinad, Tamil Nadu-India.

²Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu -Jammu-India

Citation: Bharathiraja and Lalit Upadhyay (2022). An Overview of Mycorrhiza in Pines: Research, Species, and Applications. *Plant Science Archives*. 11-12. DOI: <https://doi.org/10.5147/PSA.2022.7.4.11>

Corresponding Author: Lalit Upadhyay | E-Mail: (lupadhyay@gmail.com)

Received 09 August 2022 | Revised 22 September 2022 | Accepted 26 October 2022 | Available Online November 27 2022

ABSTRACT

Mycorrhizal associations are a fundamental aspect of pine tree ecology and biology, playing a crucial role in nutrient uptake, stress resistance, and overall plant health. This review aims to provide a comprehensive overview of the current research on mycorrhizal associations in pine species, highlighting the diversity of mycorrhizal fungi involved, and exploring the practical applications of these symbiotic relationships in forestry and conservation. We discuss the mechanisms of mycorrhizal interactions, the ecological significance of these associations, and the potential benefits of harnessing mycorrhizal fungi for sustainable forest management and reforestation efforts.

Keywords: Mycorrhiza, Pine trees, Ectomycorrhizal fungi, Arbuscular mycorrhizal fungi, Symbiosis, Nutrient uptake, Soil health

Introduction

Pine trees (genus *Pinus*) are among the most ecologically and economically important coniferous species worldwide, dominating vast forested areas and contributing significantly to timber production, carbon sequestration, and biodiversity conservation [1-2]. The success of pine trees in various environments is largely attributed to their symbiotic relationships with mycorrhizal fungi. Mycorrhizae, which are mutualistic associations between plant roots and fungi, enhance the ability of pines to absorb water and essential nutrients, particularly phosphorus and nitrogen, from the soil. This symbiosis not only improves the growth and health of pine trees but also increases their resilience to environmental stresses such as drought, soil salinity, and pathogen attacks [3-4]. Over the past few decades, extensive research has been conducted to understand the intricate dynamics of mycorrhizal associations in pine species. This body of work has revealed the diversity of mycorrhizal fungi that form symbiotic relationships with pines, including both ectomycorrhizal (EM) and arbuscular mycorrhizal (AM) fungi. Ectomycorrhizal fungi, in particular, are predominant in pine forests, forming a sheath around the roots and extending their hyphae into the surrounding soil to facilitate nutrient uptake. Arbuscular mycorrhizal fungi, although less common in pines, also play a role in enhancing nutrient acquisition and soil structure [5-7].

Research on Mycorrhizal Associations in Pines

Research on mycorrhizal associations in pines has primarily focused on identifying the species of fungi involved and elucidating the mechanisms through which these symbiotic relationships benefit the host plants. Molecular techniques, such as DNA sequencing and metagenomics, have significantly advanced our understanding of the diversity and composition of mycorrhizal communities in pine forests. Studies have shown that the mycorrhizal fungi associated with pines are highly diverse, with species composition varying according to geographical location, soil type, and environmental conditions [8-9]. The ecological significance of mycorrhizal associations in pine ecosystems cannot be overstated.

Mycorrhizal fungi enhance nutrient cycling and soil fertility, promoting the growth and productivity of pine forests [10]. These fungi also contribute to soil aggregation and stability, which are critical for preventing soil erosion and maintaining ecosystem health. Furthermore, mycorrhizal associations play a pivotal role in the establishment and survival of pine seedlings, facilitating their growth in nutrient-poor soils and under adverse environmental conditions.

Species Diversity of Mycorrhizal Fungi in Pines

The diversity of mycorrhizal fungi associated with pine trees is immense, with hundreds of species documented in various pine-dominated ecosystems. Ectomycorrhizal fungi, including genera such as *Suillus*, *Rhizopogon*, *Cenococcum*, and *Pisolithus*, are particularly prevalent in pine forests. These fungi form a dense network of hyphae around the roots, effectively increasing the surface area for nutrient and water absorption. Arbuscular mycorrhizal fungi, though less commonly associated with pines, include genera like *Glomus* and *Acaulospora*, which penetrate the root cells and form arbuscules to facilitate nutrient exchange [11].

The composition of mycorrhizal communities in pine forests is influenced by various biotic and abiotic factors. Biotic factors include the species of pine and the presence of other plants, which can affect the fungal species composition through competitive or facilitative interactions. Abiotic factors such as soil pH, temperature, moisture, and nutrient availability also play a crucial role in determining the diversity and abundance of mycorrhizal fungi. Understanding these factors is essential for managing mycorrhizal associations in pine forests to enhance tree growth and forest productivity [12].

Applications of Mycorrhizal Fungi in Forestry and Conservation

The practical applications of mycorrhizal fungi in forestry and conservation are vast and hold significant potential for sustainable forest management. Mycorrhizal inoculation, which involves introducing specific mycorrhizal fungi to seedlings in nurseries or during planting, has been shown to enhance the

growth and survival of pine trees [13]. This practice is particularly beneficial in reforestation and afforestation projects, where it can help establish healthy and resilient pine populations in degraded or nutrient-poor soils.

Mycorrhizal fungi also have applications in phytoremediation, the use of plants and their associated microorganisms to clean up contaminated environments. Pines inoculated with specific mycorrhizal fungi can be used to remediate soils contaminated with heavy metals or other pollutants, as these fungi can enhance the tolerance and uptake of contaminants by the plants. Additionally, mycorrhizal associations can improve soil health and fertility, contributing to the restoration of degraded ecosystems and the conservation of biodiversity [14-18]. The use of mycorrhizal fungi in forestry practices also supports climate change mitigation efforts. By promoting the growth and health of pine forests, mycorrhizal associations enhance carbon sequestration, thus helping to reduce atmospheric CO₂ levels. Furthermore, mycorrhizal fungi can increase the resilience of pine forests to climate-related stresses, such as drought and extreme temperatures, ensuring the long-term sustainability of these vital ecosystems [19-22].

Conclusion

The study of mycorrhizal associations in pine trees has significantly advanced our understanding of the complex interactions between plants and fungi and their implications for forest ecology and management. The diversity of mycorrhizal fungi associated with pines and their critical role in nutrient uptake, stress resistance, and overall plant health underscore the importance of these symbiotic relationships in sustaining pine forests. The practical applications of mycorrhizal fungi in forestry and conservation offer promising solutions for enhancing tree growth, restoring degraded ecosystems, and mitigating climate change. As research continues to uncover the intricacies of mycorrhizal associations in pines, it is essential to integrate this knowledge into forest management practices and conservation strategies. By harnessing the potential of mycorrhizal fungi, we can improve the productivity and resilience of pine forests, ensuring their sustainability for future generations. Continued collaboration between researchers, foresters, and conservationists will be crucial in translating scientific insights into practical applications that benefit both pine ecosystems and the broader environment.

References

- Smith, S. E., & Read, D. J. (2008). *Mycorrhizal Symbiosis*. Academic Press.
- Brundrett, M. C. (2009). Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant and Soil*, 320(1-2), 37-77.
- Courty, P. E., Buee, M., Diedhiou, A. G., Frey-Klett, P., Le Tacon, F., Rineau, F., ... & Garbaye, J. (2010). The role of ectomycorrhizal communities in forest ecosystem processes: new perspectives and emerging concepts. *Soil Biology and Biochemistry*, 42(5), 679-698.
- Agerer, R. (2001). Exploration types of ectomycorrhizae. *Mycorrhiza*, 11(2), 107-114.
- van der Heijden, M. G., & Horton, T. R. (2009). Socialism in soil? The importance of mycorrhizal fungal networks for facilitation in natural ecosystems. *Journal of Ecology*, 97(6), 1139-1150.
- Cairney, J. W., & Meharg, A. A. (2003). Ericoid mycorrhiza: a partnership that exploits harsh edaphic conditions. *European Journal of Soil Science*, 54(4), 735-740.
- Smith, F. A., & Smith, S. E. (2011). Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annual Review of Plant Biology*, 62, 227-250.
- Garbaye, J. (1994). Helper bacteria: a new dimension to the mycorrhizal symbiosis. *New Phytologist*, 128(2), 197-210.
- Hoeksema, J. D., Chaudhary, V. B., Gehring, C. A., Johnson, N. C., Karst, J., Koide, R. T., ... & Umbanhowar, J. (2010). A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecology Letters*, 13(3), 394-407.
- Smith, S. E., Jakobsen, I., Gronlund, M., & Smith, F. A. (2011). Roles of arbuscular mycorrhizas in plant phosphorus nutrition: interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiology*, 156(3), 1050-1057.
- Johnson, N. C., Graham, J. H., & Smith, F. A. (1997). Functioning of mycorrhizal associations along the mutualism-parasitism continuum. *New Phytologist*, 135(4), 575-585.
- George, E., Marschner, H., & Jakobsen, I. (1995). Role of arbuscular mycorrhizal fungi in uptake of phosphorus and nitrogen from soil. *Critical Reviews in Biotechnology*, 15(3-4), 257-270.
- Allen, M. F., & Allen, E. B. (1986). Water relations of xeric grasses in the field: interactions of mycorrhizas and competition. *New Phytologist*, 104(4), 559-571.
- Marschner, H., & Dell, B. (1994). Nutrient uptake in mycorrhizal symbiosis. *Plant and Soil*, 159(1), 89-102.
- Anderson, I. C., & Cairney, J. W. (2004). Diversity and ecology of soil fungal communities: increased understanding through the application of molecular techniques. *Environmental Microbiology*, 6(8), 769-779.
- Parniske, M. (2008). Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nature Reviews Microbiology*, 6(10), 763-775.
- Finlay, R. D. (2008). Ecological aspects of mycorrhizal symbiosis: with special emphasis on the functional diversity of interactions involving the extraradical mycelium. *Journal of Experimental Botany*, 59(5), 1115-1126.
- Read, D. J., & Perez-Moreno, J. (2003). Mycorrhizas and nutrient cycling in ecosystems—a journey towards relevance? *New Phytologist*, 157(3), 475-492.
- Smith, S. E., & Read, D. J. (1997). *Mycorrhizal Symbiosis*. Academic Press.
- Sanders, I. R., & Fitter, A. H. (1992). Evidence for differential responses between host-fungus combinations of vesicular-arbuscular mycorrhizas from a grassland. *Mycological Research*, 96(5), 415-419.
- Allen, M. F. (1991). *The Ecology of Mycorrhizae*. Cambridge University Press.
- Buee, M., Rossignol, M., Jauneau, A., Ranjeva, R., & Bécard, G. (2000). The pre-symbiotic growth of arbuscular mycorrhizal fungi is induced by a branching factor partially purified from plant root exudates. *Molecular Plant-Microbe Interactions*, 13(6), 693-698.